

# Spherical Harmonics [lam17]

Any function  $g(\theta, \phi)$  defined on the unit sphere and expressed by polar angle  $\theta$  and azimuthal angle  $\phi$  can be expanded as a series of spherical harmonics:


$$g(\theta, \phi) = \sum_{l=0}^{\infty} \sum_{m=-l}^l C_{lm} Y_{lm}(\theta, \phi), \quad C_{lm} = \int_0^{2\pi} d\phi \int_0^{\pi} d\theta \sin \theta Y_{lm}^*(\theta, \phi) g(\theta, \phi).$$

Spherical harmonics are a complete set of orthonormal functions, composed of associated Legendre functions for  $\theta$  and harmonic oscillations for  $\phi$ :

$$Y_{lm}(\theta, \phi) \doteq \sqrt{\frac{2l+1}{4\pi} \frac{(l-m)!}{(l+m)!}} P_l^m(\cos \theta) e^{im\phi}.$$

Associated Legendre functions:

- Rodrigues' generator:  $P_l^m(u) = \frac{(-1)^m}{2^l l!} (1-u^2)^{m/2} \frac{d^{l+m}}{du^{l+m}} (u^2-1)^l$ .
- Relation to polynomials:  $P_l^m(u) = (-1)^m (1-u^2)^{m/2} \frac{d^m}{du^m} P_l(u)$ .
- Orthogonality:  $\int_{-1}^{+1} du P_l^m(u) P_l^m(u) = \frac{2}{2l+1} \frac{(l+m)!}{(l-m)!} \delta_{ll}$ .
- Relation:  $P_l^{-m}(u) = (-1)^m \frac{(l-m)!}{(l+m)!} P_l^m(u)$ .
- Case  $m=0$ :  $P_l^0(x) = P_l(x)$ .
- Case  $l=1$ :  $P_1^1(x) = -\sqrt{1-x^2}$ ,  $P_1^0(x) = x$ ,  $P_1^{-1}(x) = \frac{1}{2}\sqrt{1-x^2}$ .



- Case  $l=2$ :  $P_2^2(x) = 3(1-x^2)$ ,  $P_2^1(x) = -3x\sqrt{1-x^2}$ ,  
 $P_2^0(x) = \frac{1}{2}(3x^2-1)$ ,  $P_2^{-1}(x) = \frac{1}{2}x\sqrt{1-x^2}$ ,  
 $P_2^{-2}(x) = \frac{1}{8}(1-x^2)$ .

Attributes of spherical harmonics:

- Complex conjugate function:  $Y_{lm}^*(\theta, \phi) = (-1)^m Y_{l, -m}(\theta, \phi)$ .
- Orthonormality:  $\int_0^{2\pi} d\phi \int_0^\pi d\theta \sin \theta Y_{l'm'}^*(\theta, \phi) Y_{lm}(\theta, \phi) = \delta_{l'l} \delta_{m'm}$ .
- Completeness:  $\sum_{l=0}^{\infty} \sum_{m=-l}^l Y_{lm}^*(\theta', \phi') Y_{lm}(\theta, \phi) = \delta(\phi - \phi') \delta(\cos \theta - \cos \theta')$ .
- Case  $l = 0$ :  $Y_{00} = \frac{1}{\sqrt{4\pi}}$ .
- Case  $l = 1$ :  $Y_{11} = -\sqrt{\frac{3}{8\pi}} \sin \theta e^{i\phi}$ ,  $Y_{10} = \sqrt{\frac{3}{4\pi}} \cos \theta$ ,  $Y_{1,-1} = \sqrt{\frac{3}{8\pi}} \sin \theta e^{-i\phi}$ .
- Case  $l = 2$ :  $Y_{22} = \sqrt{\frac{15}{32\pi}} \sin^2 \theta e^{2i\phi}$ ,  $Y_{21} = -\sqrt{\frac{15}{8\pi}} \sin \theta \cos \theta e^{i\phi}$ ,  
 $Y_{20} = \sqrt{\frac{5}{4\pi}} \left(\frac{3}{2} \cos^2 \theta - \frac{1}{2}\right)$ ,  
 $Y_{2,-1} = \sqrt{\frac{15}{8\pi}} \sin \theta \cos \theta e^{-i\phi}$ ,  $Y_{2,-2} = \sqrt{\frac{15}{32\pi}} \sin^2 \theta e^{-2i\phi}$ .

Addition theorem for spherical harmonics:

$$P_l(\cos \gamma) = \frac{4\pi}{2l+1} \sum_{m=-l}^l Y_{lm}^*(\theta', \phi') Y_{lm}(\theta, \phi),$$

where  $\gamma$  is the angle between position vector  $\mathbf{x}$  with angular coordinates  $\theta, \phi$  and position vector  $\mathbf{x}'$  with angular coordinates  $\theta', \phi'$ . The angles are thus related as follows:  $\cos \gamma = \cos \theta \cos \theta' + \sin \theta \sin \theta' \cos(\phi - \phi')$ .

Sum rule emerging in the limit  $\gamma \rightarrow 0$ :

$$\sum_{m=-l}^l |Y_{lm}(\theta, \phi)|^2 = \frac{2l+1}{4\pi}.$$

Application to electrostatic potential of point charge:

$$\frac{1}{|\mathbf{x} - \mathbf{x}'|} = \sum_{l=0}^{\infty} \frac{r_{<}^l}{r_{>}^{l+1}} P_l(\cos \gamma) = \sum_{l=0}^{\infty} \sum_{m=-l}^l \frac{4\pi}{2l+1} \frac{r_{<}^l}{r_{>}^{l+1}} Y_{lm}^*(\theta', \phi') Y_{lm}(\theta, \phi),$$

where  $r_{<}$  and  $r_{>}$  are the magnitudes of the shorter and longer vectors  $\mathbf{x}$  and  $\mathbf{x}'$ , respectively.