
## Condensation and evaporation [tln32]

Supersaturated gases and superheated liquids owe their metastable existence to the surface tension  $\sigma$ . Consider a liquid droplet in equilibrium with the surrounding vapor, implying  $T_l = T_g$ ,  $\mu_l = \mu_g$ , and  $p_l > p_g$  because of surface tension.

For a vapor bubble surrounded by liquid, the argument proceeds along analogous lines.



Work done if droplet expands or contracts:  $\delta W = -p_l dV_l - p_g dV_g + \sigma dA$ . Grand potential:  $\Omega(T, V, \mu) = -p_l V_l - p_g V_g + \sigma A$ .

$$\Rightarrow \ \Omega(T, V, \mu) = -\frac{4\pi}{3}R^3p_l - \left(V_{tot} - \frac{4\pi}{3}R^3\right)p_g + 4\pi R^2\sigma.$$

Equilibrium condition:  $(\partial\Omega/\partial R)_{T,V,\mu} = 0 \Rightarrow 4\pi R^2 (p_g - p_l) + 8\pi R\sigma = 0.$ Excess pressure in droplet:  $p_l - p_g = 2\sigma/R.$ Gibbs-Duhem equations (with dT = 0),  $N_l d\mu_l = V_l dp_l$ ,  $N_g d\mu_g = V_g dp_g.$ Chemical equilibrium:  $d\mu_l = d\mu_g \Rightarrow (V_l/N_l) dp_l = (V_g/N_g) dp_g.$ Differential excess pressure:  $d(p_l - p_g) = \frac{V_g/N_g - V_l/N_l}{V_l/N_l} dp_g = d\left(\frac{2\sigma}{R}\right).$ Use  $\frac{V_g}{N_g} \gg \frac{N_l}{V_l}, \ \frac{V_g}{N_g} \simeq \frac{k_B T}{p_g} \Rightarrow \frac{k_b T/p_g}{V_l/N_l} dp_g = d\left(\frac{2\sigma}{R}\right).$ Integrate  $\frac{dp_g}{p_g} = \frac{V_l}{N_l k_B T} d\left(\frac{2\sigma}{R}\right)$  from  $\infty$  to R.  $\Rightarrow \ln \frac{p_g(R)}{p_g(\infty)} = \frac{2\sigma V_l}{RN_l k_B T} = \frac{2\sigma m}{R\rho_l k_B T} \Rightarrow p_g(R) = p_g(\infty) \exp\left(\frac{2\sigma m}{R\rho_l k_B T}\right).$ Only liquid droplets of a particular radius  $R_c$  coexist with the supersaturated

Only liquid droplets of a particular radius  $R_c$  coexist with the supersaturated gas phase. Droplets with  $R < R_c$  will shrink. Droplets with  $R > R_c$  will grow. Hence the condensation process at pressure  $p = p_g(R_c)$  can be initiated by the presence of droplets with radius  $R > R_c$ .

Metastability depends on the absence of droplets with radius  $R > R_c$ . The boundary of the metastable region (spinodal line) corresponds to a value of  $R_c$  comparable to the molecular radius. Supersaturation cannot be pushed beyond that point.